

ELECTROMAGNETIC FORM FACTORS OF BARYONS

R. BIJKER

*Instituto de Ciencias Nucleares,
Universidad Nacional Autónoma de México,
A.P. 70-543, 04510 México D.F., México*

A. LEVIATAN

*Racah Institute of Physics,
The Hebrew University, Jerusalem 91904, Israel*

Nucleon elastic and transition form factors are analyzed in the framework of a collective model of baryons. Effects of meson cloud couplings and relativistic corrections are considered.

1 Introduction

Electromagnetic form factors of the nucleon and its excitations (baryon resonances) provide a powerful tool to investigate the structure of the nucleon. These form factors can be measured in electroproduction as a function of the four-momentum squared $q^2 = -Q^2$ of the virtual photon.

In this contribution we present a simultaneous study of the elastic form factors of the nucleon and the transition form factors, for both of which there exists exciting new data¹. The analysis is carried out in the framework of a collective model of the nucleon². We address the effects of relativistic corrections and couplings to the meson cloud.

2 Collective model of baryons

In the recently introduced collective model of baryons² the radial excitations of the nucleon are described as vibrational and rotational excitations of an oblate top. The electromagnetic form factors are obtained by folding with a distribution of the charge and magnetization over the entire volume of the baryon. All calculations are carried out in the Breit frame.

2.1 Elastic form factors

In the collective model, the electric and magnetic form factors of the nucleon are expressed in terms of a common intrinsic dipole form factor². The effects of the meson cloud surrounding the nucleon are taken into account by coupling the nucleon form factors to the isoscalar vector mesons ω and ϕ and the isovector vector meson ρ . The coupling is carried out at the level of the Sachs form

Table 1: Ratios of helicity amplitudes.

N(1535) S_{11}	$A_{1/2}^n/A_{1/2}^p$	-0.84 ± 0.15^8	-0.81^2	-0.90
N(1520) D_{13}	$A_{3/2}^p/A_{1/2}^p$	$-2.5 \pm 0.2 \pm 0.4^9$	-2.53^2	-2.66

factors. The large width of the ρ meson is taken into account according to the prescription of³, and the constraints from perturbative QCD are imposed by scaling Q^2 with the strong coupling constant⁴.

We carried out a simultaneous fit to all four electromagnetic form factors of the nucleon and the nucleon charge radii, and found good agreement with the world data, including the new data for the neutron form factors (square boxes in Fig. 1). The oscillations around the dipole values are due to the meson cloud couplings.

2.2 Transition form factors

Recent experiments on eta-photoproduction have yielded valuable new information on the helicity amplitudes of the N(1520) D_{13} and N(1535) S_{11} resonances. In Table 1 we show the model-independent ratios of photocouplings that have been extracted from the new data^{8,9}. These values are in excellent agreement with those of the collective model². In the last column we show the effect of relativistic corrections to the electromagnetic transition operator. In Fig. 2 we show the N(1535) S_{11} proton helicity amplitude. The new data are indicated by diamonds and square boxes. The effect of relativistic corrections (dashed line) to the nonrelativistic results (solid line) in the collective model of² is small, and shows up mainly at small values of Q^2 .

3 Summary and conclusions

We presented a simultaneous analysis of the four elastic form factors of the nucleon and the transition form factors in a collective model of baryons, and found good agreement with the new data presented at this conference.

The deviations of the nucleon form factors from the dipole form were attributed to couplings to the meson cloud which were taken into account using vector meson dominance. The helicity amplitudes of the N(1520) D_{13} and N(1535) S_{11} resonances as well as their model-independent ratios are reproduced well. The effect of relativistic corrections is small, and shows up




Figure 1: Nucleon form factors as a function of Q^2 ($\text{GeV}/c)^2$: a) G_E^p/F_D from ⁵, b) $G_M^p/\mu_p F_D$ from ⁵, c) G_E^n from ⁶ and d) $G_M^n/\mu_n F_D$ from ⁷; IJL from ³ and GK from ⁴; $F_D = 1/(1 + Q^2/0.71)^2$.

mainly at small values of Q^2 .

In conclusion, the present analysis of electromagnetic couplings shows that the collective model of baryons provides a good overall description of the available data.




Figure 2: $N(1535)S_{11}$ proton helicity amplitude in $10^{-3} \text{ GeV}^{-1/2}$ as a function of Q^2 ($\text{GeV}/c)^2$. A factor of $+i$ is suppressed. The data are taken from¹⁰.

Acknowledgements

This work is supported in part by DGAPA-UNAM under project IN101997 and by grant No. 94-00059 from the United States-Israel Binational Science Foundation (BSF).

References

1. See e.g. D. Drechsel, D. Rebreyend and P. Stoler, these proceedings.
2. R. Bijker, F. Iachello and A. Leviatan, *Ann. Phys. (N.Y.)* **236**, 69 (1994); *Phys. Rev. C* **54**, 1935 (1996).
3. F. Iachello, A.D. Jackson and A. Lande, *Phys. Lett. B* **43**, 191 (1973).
4. M. Gari and W. Krümpelmann, *Z. Phys. A* **322**, 689 (1985).

5. A.F. Sill et al., *Phys. Rev. D* **48**, 29 (1993); R.C. Walker et al., *Phys. Rev. D* **49**, 5671 (1994).
6. C.E. Jones-Woodward et al., *Phys. Rev. C* **44**, R571 (1991); A.K. Thompson et al., *Phys. Rev. Lett.* **68**, 2901 (1992); M. Meyerhoff et al., *Phys. Lett. B* **327**, 201 (1994); T. Eden et al., *Phys. Rev. C* **50**, R1749 (1994); H. Schmieden and P. Grabmayr, these proceedings.
7. R.G. Arnold et al., *Phys. Rev. Lett.* **61**, 806 (1988); S. Rock et al., *Phys. Rev. D* **46**, 24 (1992); A. Lung et al., *Phys. Rev. Lett.* **70**, 718 (1993); P. Markowitz et al., *Phys. Rev. C* **48**, R5 (1993); H. Gao et al., *Phys. Rev. C* **50**, R546 (1994); H. Anklin et al., *Phys. Lett. B* **336**, 313 (1994); E.E.W. Bruins et al., *Phys. Rev. Lett.* **75**, 21 (1995); H. Anklin et al., *Phys. Lett. B* **428**, 248 (1998).
8. N.C. Mukhopadhyay, J.-F. Zhang and M. Benmerrouche, *Phys. Lett. B* **364**, 1 (1995).
9. N.C. Mukhopadhyay and N. Mathur, preprint nucl-th/9807003.
10. V.D. Burkert, *Int. J. Mod. Phys. E* **1**, 421 (1992); B. Krusche et al., *Phys. Lett. B* **397**, 171 (1997); C.S. Armstrong et al., nucl-ex/9811001.